

## Now that you've had your **ARESTIN** treatment, what comes next?



### AVOID

- Hard, crunchy, or sticky foods for 1 week
- Touching any treated areas
- Devices designed to clean around the treated site (like floss and toothpicks) for 10 days




### EXPECT

- Mild-to-moderate sensitivity in the first week



### TELL

- Your dentist or hygienist if you're experiencing any pain, itching, or swelling, or if possible signs and symptoms of hypersensitivity occur



*Ask your dental professional to mark infected areas treated with ARESTIN.*

### NOTES:

---

---

---

### YOUR NEXT APPOINTMENT:

Date 

---

Time 

---

### INDICATION

ARESTIN® (minocycline HCl) Microspheres, 1mg is used in combination with scaling and root planing (SRP) procedures to treat patients with adult periodontitis (gum disease). ARESTIN® may be used as part of an overall oral health program that includes good brushing and flossing habits and SRP.

### IMPORTANT SAFETY INFORMATION

- Do not take ARESTIN if you are allergic to minocycline or tetracyclines. Ask your dentist or pharmacist for a list of these drugs if you are not sure. Swelling of the face, itching of the skin, fever and enlarged lymph nodes have been reported with the use of ARESTIN. Some of these reactions were serious. Tell your dentist right away if you have any signs of allergic reaction, such as skin reactions or trouble breathing, or if you have an exaggerated sunburn reaction.

Please see additional Important Safety Information on reverse and accompanying full Prescribing Information.

## QUESTIONS?

Be sure to ask your dentist or hygienist.

**IMPORTANT SAFETY INFORMATION (CONT'D)**

- THE USE OF TETRACYCLINE CLASS DRUGS, INCLUDING ARESTIN, DURING TOOTH DEVELOPMENT MAY CAUSE PERMANENT DISCOLORATION OF THE TEETH, AND THEREFORE SHOULD NOT BE USED IN CHILDREN OR IN PREGNANT OR NURSING WOMEN.
- ARESTIN has not been studied in patients with weakened immune systems (such as patients with HIV infections or diabetes, or those receiving chemotherapy or radiation). Tetracyclines, including oral minocycline, have been associated with the development of autoimmune syndrome with symptoms such as joint pain, muscle pain, rash, swelling, fever, enlarged lymph nodes, and general body weakness. Tell your doctor about any health problems you have, including whether you have had oral candidiasis ("thrush") in the past, and about all medications you are taking.
- In clinical studies, the most frequently reported non-dental side effects were headache, infection, flu symptoms, and pain.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch) or call 1-800-FDA-1088.

Please see additional Important Safety Information on reverse and accompanying full Prescribing Information